



## Canadian Water Quality Guidelines for the Protection of Aquatic Life

## CHLORINATED ETHENES 1,1,2-trichloroethene (trichloroethylene)

**T**richloroethylene ( $C_2HCl_3$ ; CAS 79-01-06) and TCE are common names for 1,1,2-trichloroethene; a colourless liquid with high volatility (vapour pressure = 8 kPa at 20°C) and moderate water solubility ( $1.1 \text{ g}\cdot\text{L}^{-1}$  at 20°C). TCE is a degreasing solvent in automotive and metal industries, and is used in dry cleaning facilities and various consumer products (Hughes et al. 1994). Canadian production of TCE ceased in 1985.

TCE enters the aquatic environment through industrial discharges, landfill leaching, accidental spills, and improper storage and disposal. Atmospheric transport and deposition of TCE is a significant route of entry to aquatic environments in nonindustrial areas (USEPA 1979). In the 1980s, surface water concentrations in Canada generally did not exceed  $1 \text{ }\mu\text{g}\cdot\text{L}^{-1}$ , though site-specific residues up to  $90 \text{ }\mu\text{g}\cdot\text{L}^{-1}$  were reported (Kaiser and Comba 1983; 1986; Lum and Kaiser 1986). TCE concentrations in groundwater are generally higher due to minimal volatilization (Zoeteman et al. 1980). Residues from  $0.06\text{--}425 \text{ mg}\cdot\text{L}^{-1}$  have been measured near industrial sites in British Columbia and Manitoba, respectively (Golder Associates 1989; UMA Engineering Ltd. 1992).

The low  $\log K_{ow}$  (2.29) of TCE suggests that it will not bioconcentrate to a high degree. BCFs for freshwater organisms vary from 17 to 1160 for bluegill sunfish (*Lepomis macrochirus*) and green alga (*Chlorella fusca vacuolata*), respectively (Barrows et al. 1980; Geyer et al. 1984). Further TCE residues in several marine species are reportedly  $\leq 30.0 \text{ mg}\cdot\text{kg}^{-1}$  ww (Pearson and McConnell 1975; Ofstad et al. 1981; Wang et al. 1985).

Volatilization is the principal removal process for TCE from the aquatic environment. Laboratory studies have indicated half-lives from 17.7 min to 3.4 d, depending on the rate of mixing (Jensen and Rosenberg 1975; Dilling 1977; Peng et al. 1994). Estimated half-lives in shallow ponds, lakes, and rivers are  $<12$  d (Smith et al. 1980; Lay et al. 1984). TCE may be more persistent in marine systems with estimated half-lives of 13–28 d (Wakeham et al. 1983). Freshwater rich in organic matter may lose up to 40% of dissolved TCE residues through sorption (Dilling et al. 1975). Limited microbial degradation occurs under anoxic conditions (Barrio-Lage et al. 1988).

Despite rapid dissipation rates, TCE may have long-term ecological effects. For example, in a natural pond community with an initial TCE concentration of  $25.0 \text{ mg}\cdot\text{L}^{-1}$ , zooplankton (*Daphnia* sp.) abundance and phytoplankton richness and abundance remained low 43 d after application, although 98% of the TCE had disappeared within 15 d (Lay et al. 1984).

### Water Quality Guideline Derivation

The interim Canadian water quality guideline for 1,1,2-trichloroethene for the protection of freshwater life was developed based on the CCME protocol (CCME 1991a).

### Freshwater Life

Acute toxicity (48- to 96-h  $LC_{50}$ ) values for TCE vary from  $28.2$  to  $213 \text{ mg}\cdot\text{L}^{-1}$  for juvenile American flagfish (*Jordanella floridae*) and golden orfes (*Leuciscus idus*), respectively (Slooff et al. 1983; ATRG 1988). Fathead minnows (*Pimephales promelas*) exposed for 96 h to  $21.9 \text{ mg}\cdot\text{L}^{-1}$  TCE experienced narcosis, gill swelling, melanization, and loss of equilibrium (Alexander et al. 1978). Following exposure to  $5.0 \text{ mg}\cdot\text{L}^{-1}$  of TCE for 24 h, rainbow trout (*Oncorhynchus mykiss*) increased their respiration rate (Slooff 1979). Chronic toxicity data indicate that brook trout (*Salvelinus fontinalis*) experience a decrease in swim-up survival and 120-d fry weight and growth after exposure to  $0.21 \text{ mg}\cdot\text{L}^{-1}$  (ATRG 1988). Flagfish fry suffer 100% mortality after exposure to  $20.9 \text{ mg}\cdot\text{L}^{-1}$  TCE for 28 d (ATRG 1988).

Water fleas (*Daphnia magna*) are the most sensitive aquatic invertebrate to TCE with a 48-h  $EC_{50}$  for

**Table 1. Water quality guidelines for 1,1,2-trichloroethene for the protection of aquatic life (CCME 1991b).**


Aquatic life	Guideline value ( $\mu\text{g}\cdot\text{L}^{-1}$ )
Freshwater	21*
Marine	NRG <sup>†</sup>

\*Interim guideline.

<sup>†</sup>No recommended guideline.

immobilization of 7.76 mg·L<sup>-1</sup> (Abernethy et al. 1986). Other studies report 48-h LC<sub>50</sub>s from 48.0 to 75.0 mg·L<sup>-1</sup> for mosquitoes (*Aedes aegypti*) and polyps (*Hydra oligactis*), respectively (Slooff et al. 1983). Some phytoplankton species are quite tolerant of TCE. For example, *Selenastrum capricornutum* has a 96-h NOEC for growth of 175 mg·L<sup>-1</sup>, and *Scenedesmus pannonicus* has a 192-h NOEC of >1000 mg·L<sup>-1</sup> TCE (Slooff et al. 1983).

The interim water quality guideline for TCE for the protection of freshwater life is 21 µg·L<sup>-1</sup>. It was derived by multiplying the LOEC of 0.21 mg·L<sup>-1</sup> for brook trout (ATRG 1988) by a safety factor of 0.1 (CCME 1991a, 1991b).



**Figure 1. Select freshwater toxicity data for 1,1,2-trichloroethene.**

**References**

Abernethy, S., A.M. Bobra, W.Y. Shiu, R.G. Wells, and D. Mackay. 1986. Acute lethal toxicity of hydrocarbons and chlorinated hydrocarbons to two planktonic crustaceans: The key role of organism-water partitioning. *Aquat. Toxicol.* 8:163-174.

Alexander, H.C., W.M. McCarty, and E.A. Bartlett. 1978. Toxicity of perchloroethylene, trichloroethylene, 1,1,1-trichloroethane, and methylene chloride to fathead minnows. *Bull. Environ. Contam. Toxicol.* 20:344-352.

ATRG (Aquatic Toxicity Research Group). 1988. Aquatic toxicity of multiple organic compounds. II. Chlorinated ethanes and chlorinated ethylenes. A summary report prepared for the Ontario Ministry of the

Environment by the Aquatic Toxicity Research Group, Lakehead University, Thunder Bay, ON.

Barrio-Lage, G.A., F.Z. Parsons, and P.A. Lorenzo. 1988. Inhibition and stimulation of trichloroethylene biodegradation in microaerophilic microcosms. *Environ. Toxicol. Chem.* 7:889-895.

Barrows, M.E., S.R. Petrocelli, K.J. Macek, and J.J. Carroll. 1980. Bio-concentration and elimination of selected water pollutants by bluegill sunfish (*Lepomis macrochirus*). In: *Dynamics, exposure and hazard assessment of toxic chemicals*, R. Haque, ed. Ann Arbor Science Publishers, Ann Arbor, MI.

CCME (Canadian Council of Ministers of the Environment). 1991a. Appendix IX—A protocol for the derivation of water quality guidelines for the protection of aquatic life (April 1991). In: *Canadian water quality guidelines*, Canadian Council of Resource and Environment Ministers. 1987. Prepared by the Task Force on Water Quality Guidelines. [Updated and reprinted with minor revisions and editorial changes in Canadian environmental quality guidelines, Chapter 4, Canadian Council of Ministers of the Environment, 1999, Winnipeg.]

———. 1991b. Appendix VII—Canadian water quality guidelines: Updates (April 1991), trichloroethylene, polychlorinated biphenyls (PCBs), and chlorinated ethanes. In: *Canadian water quality guidelines*, Canadian Council of Resource and Environment Ministers. 1987. Prepared by the Task Force on Water Quality Guidelines.

Dilling, W.L. 1977. Interphase transfer processes. II. Evaporation rates of chloromethanes, ethanes, ethylenes, propanes, and propylenes from dilute aqueous solutions. *Comparisons with theoretical predictions.* *Environ. Sci. Technol.* 11:405-409.

Dilling, W.L., N.B. Tefertiller, and G.J. Kallos. 1975. Evaporation rates and reactivities of methylene chloride, chloroform, 1,1,1-trichloroethane, trichloroethylene, tetrachloroethylene, and other chlorinated compounds in dilute aqueous solutions. *Environ. Sci. Technol.* 9:833-838.

Geyer, H., G. Politzki, and D. Freitag. 1984. Prediction of ecotoxicological behaviour of chemicals: Relationship between n-octanol-water partition coefficient and bioaccumulation of organic chemicals by alga *Chlorella*. *Chemosphere* 13:269-284.

Golder Associates. 1989. Interim report on supplementary hydrogeological investigations at 800 Terminal Ave., Vancouver, BC. Prepared by Golder Associates Ltd., Burnaby, BC. Unpub.

Hughes, K., M.E. Meek, and W. Windle. 1994. Trichloroethylene: Evaluation of risks to health from environmental exposure in Canada. *J. Environ. Health. Sci.* C12:527-543.

Jensen, S., and R. Rosenberg. 1975. Degradability of some chlorinated aliphatic hydrocarbons in seawater and sterilized water. *Water Res.* 9:659-661.

Kaiser, K.L.E., and M.E. Comba. 1983. Volatile contaminants in the Welland River watershed. *J. Gt. Lakes Res.* 9:274-280.

———. 1986. Tracking river plumes with volatile halocarbon contaminants: The St. Clair River-Lake St. Clair example. *Environ. Toxicol. Chem.* 5:965-976.

Lay, J.P., W. Schauerte, and W. Klein. 1984. Effects of trichloroethylene on the population dynamics of phyto- and zooplankton in compartments of a natural pond. *Environ. Pollut. Ser. A. Ecol. Biol.* 33:75-91.

Lum, K.R., and K.L.E. Kaiser. 1986. Organic and inorganic contaminants in the St. Lawrence River: Some preliminary results on their distribution. *Water Pollut. Res. J. Can.* 21:592-603.

Ofstad, E.B., H. Drangsholt, and G. Carlborg. 1981. Analysis of volatile halogenated organic compounds in fish. *Sci. Total Environ.* 20:205-215.

Pearson, C.R., and G. McConnell. 1975. Chlorinated C1 and C2 hydrocarbons in the marine environment. *Proc. R. Soc. London Ser. B189:305-332.*

- Peng, J., J.K. Bewtra, and N. Biswas. 1994. Volatilization of selected organic compounds from quiescent water. *J. Environ. Eng.* 120:662–669.
- Slooff, W. 1979. Detection limits of a biological monitoring system based on fish respiration. *Bull. Environ. Contam. Toxicol.* 23:517–523.
- Slooff, W., J.H. Canton, and J.L.M. Hermens. 1983. Comparison of the susceptibility of 22 freshwater species to 15 chemical compounds. I. (Sub)Acute toxicity tests. *Aquat. Toxicol.* 4:113–128.
- Smith, J.H., D.C. Bomberger Jr., and D.L. Hayes. 1980. Prediction of the volatilization rates of high-volatility chemicals from natural water bodies. *Environ. Sci. Technol.* 14:1332–1337.
- UMA Engineering Ltd. 1992. Bristol Aerospace Limited: Phase I, Groundwater contamination investigation, April 1992. Phase II, Soil investigation, September 1992. Prepared by UMA Engineering Ltd., Winnipeg.
- USEPA (U.S. Environmental Protection Agency). 1979. Water-related environmental fate of 129 priority pollutants. II. Halogenated aliphatic hydrocarbons, halogenated ethers, monocyclic aromatics, phthalate esters, polycyclic aromatic hydrocarbons, nitrosamines, and miscellaneous compounds. EPA 440/4 79-029b. USEPA, Washington, DC.
- Wakeham, S.G., A.C. Davis, and J.L. Karas. 1983. Mesocosm experiments to determine the fate and persistence of volatile organic compounds in coastal seawater. *Environ. Sci. Technol.* 17:611–617.
- Wang, T.C., R. Lenahan, M. Kanik, and J. TenEyck. 1985. The removal of trichloroethylene contaminated ground water at Vero Beach, Florida. *Arch. Environ. Contam. Toxicol.* 14:719–723.
- Zoeteman, B.C.J., K. Harmsen, J.B.H.J. Linders, C.F.H. Morra, and W. Slooff. 1980. Persistent organic pollutants in river water and groundwater of the Netherlands. *Chemosphere* 9:231–249.

Reference listing:

Canadian Council of Ministers of the Environment. 1999. Canadian water quality guidelines for the protection of aquatic life: Chlorinated ethenes—1,1,2-Trichloroethene (trichloroethylene). In: Canadian environmental quality guidelines, 1999, Canadian Council of Ministers of the Environment, Winnipeg.

For further scientific information, contact:

Environment Canada  
Guidelines and Standards Division  
351 St. Joseph Blvd.  
Hull, QC K1A 0H3  
Phone: (819) 953-1550  
Facsimile: (819) 953-0461  
E-mail: ceqg-rcqe@ec.gc.ca  
Internet: <http://www.ec.gc.ca>

For additional copies, contact:

CCME Documents  
c/o Manitoba Statutory Publications  
200 Vaughan St.  
Winnipeg, MB R3C 1T5  
Phone: (204) 945-4664  
Facsimile: (204) 945-7172  
E-mail: [spcme@chc.gov.mb.ca](mailto:spcme@chc.gov.mb.ca)